


## CD14 Antibody [clone LPSR/7738] (V5447)


| Catalog No.    | Formulation   | Size   |
|----------------|---|--------|
| V5447-100UG    | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V5447-20UG     | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug  |
| V5447SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free                          | 100 ug |

[Bulk quote request](#)


|                           |  |
|---------------------------|--|
| <b>Availability</b>       | 1-3 business days                                      |
| <b>Species Reactivity</b> | Human  |
| <b>Format</b>             | Purified   |
| <b>Host</b>               | Mouse  |
| <b>Clonality</b>          | Monoclonal (mouse origin)                              |
| <b>Isotype</b>            | Mouse IgG2, kappa                                      |
| <b>Clone Name</b>         | LPSR/7738  |
| <b>Purity</b>             | Protein A/G affinity                                   |
| <b>UniProt</b>            | P08571   |
| <b>Localization</b>       | Cell membrane  |
| <b>Applications</b>       | Immunohistochemistry (FFPE) : 1-2ug/ml                 |
| <b>Limitations</b>        | This CD14 antibody is available for research use only. |



IHC staining of FFPE human tonsil tissue with CD14 antibody (clone LPSR/7738). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human lymph node tissue with CD14 antibody (clone LPSR/7738).  
HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free CD14 antibody (clone LPSR/7738) as confirmation of integrity and purity.

## Description

Recognizes a protein of 55kDa, identified as CD14 (also known lipopolysaccharide receptor). CD14 is expressed strongly on monocytes and macrophage and weakly on the surface of neutrophils. CD14 is anchored to cells by linkage to glycosylphosphatidylinositol (GPI) and functions as a high affinity receptor for complexes of LPS and LPS binding protein (LBP). Soluble CD14, also binding to LPS, acts at physiological concentration as an LPS agonist and has, at higher concentrations, an LPS antagonizing effect in cell activation.

## Application Notes

Optimal dilution of the CD14 antibody should be determined by the researcher.

## Immunogen

A recombinant fragment of human CD14 protein (within amino acids 1-200) was used as the immunogen for the CD14 antibody.

## Storage

Aliquot the CD14 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.