
CD14 Antibody CDLA14-2 / Mouse Monoclonal [clone CDLA14-2] (V3764)


Catalog No.	Formulation	Size
V3764-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3764-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3764SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	CDLA14-2
Purity	Protein G affinity chromatography
UniProt	P08571
Localization	Cell surface, Secreted, Cytoplasmic (Golgi)
Applications	IHC (FFPE): 1-2ug/ml for 30 min at RT Western blot: 1-2ug/ml
Limitations	This CD14 Antibody CDLA14-2 is available for research use only.



IHC testing of FFPE human lymph node with CD14 Antibody CDLA14-2. HIER: boil tissue sections in 10mM Tris with 1mM EDTA, pH 9, for 10-20 min followed by cooling at RT for 20 min.



Western blot testing of human A549 cell lysate with CD14 antibody (clone LPSR/2386). Expected molecular weight: 40-55 kDa depending on glycosylation level.

Description

Coreceptor for bacterial lipopolysaccharide. In concert with LBP, binds to monomeric lipopolysaccharide and delivers it to the LY96/TLR4 complex, thereby mediating the innate immune response to bacterial lipopolysaccharide (LPS). Acts via MyD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response. Acts as a coreceptor for TLR2:TLR6 heterodimer in response to diacylated lipopeptides and for TLR2:TLR1 heterodimer in response to triacylated lipopeptides, these clusters trigger signaling from the cell surface and subsequently are targeted to the Golgi in a lipid-raft dependent pathway. [UniProt]

For an alternative antibody targeting the same monocyte and macrophage surface marker, see our CD14 Antibody LPSR/2385, a mouse monoclonal antibody featuring IHC, western blot, flow cytometry, and protein microarray validation.

Application Notes

The stated application concentrations are suggested starting points. Titration of the CD14 Antibody CDLA14-2 may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 25-148 was used as immunogen for this CD14 Antibody CDLA14-2.

Storage

Store the CD14 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Ordering: Phone: 858.663.9055 | Fax: 1.267.821.0800 | Email: info@nsjbio.com
Copyright © NSJ Bioreagents. All rights reserved.