

CD16 Antibody [clone CB16] (V5695)

Catalog No.	Formulation	Size
V5695-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5695-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5695SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	CB16
Purity	Protein G affinity
UniProt	P08637
Localization	Cell membrane, Secreted
Applications	Flow Cytometry : 1-2ug/million cells
Limitations	This CD16 antibody is available for research use only.



Description

It recognizes CD16 (FcγR3), the low-affinity receptor for IgG with an apparent molecular weight of 50-80kDa. CD16 is represented by two similar genes, CD16A (FcγR3A), which exists as a hetero-oligomeric polypeptide-

anchored form in macrophages and NK cells and CD16B (FcγRIIIB), which exist as a monomeric GPI-anchored form in neutrophils. Furthermore, there are two known polymorphisms of CD16B, NA-1 and NA-2. Individuals homozygous for NA-2 show a lower phagocytic capacity compared with NA-1. CD16 binds IgG in the form of immune complexes and shows preferential binding of IgG1 and IgG3 isotypes and minimal binding of IgG2 and IgG4. Upon IgG binding, both CD16 isoforms initiate signal transduction cascades that lead to a variety of responses including antibody-dependent cell-mediated cytotoxicity (ADCC), phagocytosis, degranulation and proliferation.

Researchers studying Fc-mediated immune signaling, natural killer cell biology, and antibody-dependent cellular cytotoxicity may also benefit from the [CD16 Antibody / Fc Gamma Receptor III Immune Marker page](#) featuring immunohistochemistry, immunofluorescence, and flow cytometry validation data for endogenous FCGR3 detection.

Application Notes

Optimal dilution of the CD16 antibody should be determined by the researcher.

Immunogen

PBLs from a NK-leukemia patient were used as the immunogen for the was used as the immunogen for the CD16 antibody.

Storage

Aliquot the CD16 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.