

CD1b Antibody [clone 100-1A5] (V2950)

Catalog No.	Formulation	Size
V2950-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2950-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2950SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgM, kappa
Clone Name	100-1A5
Purity	PEG precipitation
UniProt	P29016
Localization	Cell surface, cytoplasmic
Applications	Flow Cytometry: 0.5-1ug/10^6 cells Immunofluorescence: 0.5-1ug/ml Functional Studies (order BSA/sodium Azide-free Format):
Limitations	This CD1b antibody is available for research use only.



The mouse monoclonal antibody recognizes CD1b, a 44kDa type I glycoprotein associated with beta2-microglobulin (Workshop IV; Code T015). It is expressed on dendritic cells, Langerhans cells, thymocytes, and T acute lymphoblastic leukemia cells. The CD1 multigene family encodes five forms of the CD1 T-cell surface glycoprotein in human, designated CD1A, 1B, 1C, 1D and 1E. CD1, a type 1 membrane protein, has structural similarity to the MHC class I antigen and has been shown to present lipid antigens for recognition by T lymphocytes. Constitutive endocytosis of CD1B molecules and the differential sorting of MHC class II from lysosomes separate peptide- and lipid antigen-presenting molecules during dendritic cell maturation. CD1B is also expressed in interdigitating cells.

Application Notes

Optimal dilution of the CD1b antibody should be determined by the researcher.

Immunogen

Stimulated human leukocytes were used as the immunogen for the CD1b antibody.

Storage

Store the CD1b antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).