


CD2 Antibody [clone HuLy-m1] (V2951)

Catalog No.	Formulation	Size
V2951-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2951-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2951SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


 Citations (3)

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	HuLy-m1
Purity	Protein G affinity chromatography
UniProt	P06729
Localization	Cell surface
Applications	Functional Studies (order BSA/sodium Azide-free Format) : Flow Cytometry : 0.5-1ug/10 ⁶ cells Immunofluorescence : 0.5-1ug/ml
Limitations	This CD2 antibody is available for research use only.



Western blot testing of Jurkat cell lysate with CD2 antibody (clone HuLy-m1). Expected molecular weight ~47kDa.



Flow cytometry testing of human MOLT-4 cells with CD2 antibody (clone HuLy-m1); Red=isotype control, Blue= CD2 antibody.

Description

CD2 interacts through its amino-terminal domain with the extracellular domain of CD58 (also designated CD2 ligand) to mediate cell adhesion. CD2/CD58 binding can enhance antigen-specific T cell activation. CD2 is a transmembrane glycoprotein that is expressed on peripheral blood T lymphocytes, NK cells and thymocytes. CD58 is a heavily glycosylated protein with a broad tissue distribution in hematopoietic and other cells, including endothelium. Interaction between CD2 and its counter receptor LFA3 (CD58) on opposing cells optimizes immune system recognition, thereby facilitating communication between helper T lymphocytes and antigen-presenting cells, as well as between cytolytic effectors and target cells.

Application Notes

Optimal dilution of the CD2 antibody should be determined by the researcher.

Immunogen

Human thymocytes were used as the immunogen for the CD2 antibody.

Storage

Store the CD2 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).