


CD2 Antibody [clone RPA-2.10] (V8198)

Catalog No.	Formulation	Size
V8198-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8198-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8198SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	RPA-2.10
Purity	Protein G affinity chromatography
UniProt	P06729
Applications	Functional Studies (order BSA/azide-free Format) : Flow Cytometry : 1-2ug/10 ⁶ cells in 0.1ml
Limitations	This CD2 antibody is available for research use only.



SDS-PAGE analysis of purified, BSA-free CD2 antibody (clone RPA-2.10) as confirmation of integrity and purity.

Description

CD2 interacts through its amino-terminal domain with the extracellular domain of CD58 (also designated CD2 ligand) to

mediate cell adhesion. CD2/CD58 binding can enhance antigen-specific T cell activation. CD2 is a transmembrane glycoprotein that is expressed on peripheral blood T lymphocytes, NK cells and thymocytes. CD58 is a heavily glycosylated protein with a broad tissue distribution in hematopoietic and other cells, including endothelium. Interaction between CD2 and its counter receptor LFA3 (CD58) on opposing cells optimizes immune system recognition, thereby facilitating communication between helper T lymphocytes and antigen-presenting cells, as well as between cytolytic effectors and target cells.

Application Notes

Optimal dilution of the CD2 antibody should be determined by the researcher.

Immunogen

A recombinant full-length human CD2 protein was used as the immunogen for the CD2 antibody.

Storage

Store the CD2 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).