


## CD22 Antibody [clone BLCAM/1796] (V3784)


Catalog No.	Formulation	Size
V3784-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3784-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3784SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Bulk quote request


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2b, kappa
<b>Clone Name</b>	BLCAM/1796
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P20273
<b>Localization</b>	Cell surface, cytoplasmic
<b>Applications</b>	ELISA : 2-4ug/ml (order BSA/azide-free format) Western Blot : 1-2ug/ml Flow Cytometry : 1-2ug/10 <sup>6</sup> cells Immunofluorescence : 1-2ug/ml
<b>Limitations</b>	This CD22 antibody is available for research use only.




Immunofluorescent staining of human Ramos cells with CD22 antibody (clone BLCAM/1796, green) and Reddot nuclear stain (red).



Flow cytometry testing of human Ramos cells with CD22 antibody (clone BLCAM/1796);  
Red=isotype control, Blue= CD22 antibody.




Western blot testing of human Raji cell lysate with CD22 antibody (clone BLCAM/1796).  
Expected molecular weight: 76-150 kDa depending on glycosylation level.



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using CD22 antibody (clone BLCAM/1796). These results demonstrate the foremost specificity of the BLCAM/1796 mAb.

Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free CD22 antibody (clone BLCAM/1796) as confirmation of integrity and purity.

## Description

Recognizes a protein of 130-140kDa, identified as CD22 (also known as BL-CAM). CD22 expression is restricted to normal and neoplastic B cells and is absent from other haemopoietic cell types. In B-cell ontogeny, CD22 is first expressed in the cytoplasm of pro-B and pre-B cells, and on the surface as B cells mature to become IgD+. It is not expressed by plasma cells, CD22 is found highly expressed in follicular mantle and marginal zone B-cells, and while germinal center B-cells are relatively weak. CD22 is a member of the immunoglobulin superfamily and serves as an adhesion receptor for sialic acid-bearing ligands expressed on erythrocytes and all leukocyte classes. It also associates with tyrosine kinases and play a role in signal transduction and B-cell activation.

## Application Notes

Optimal dilution of the CD22 antibody should be determined by the researcher.

## Immunogen

Amino acids 52-178 from the human protein were used as the immunogen for the CD22 antibody.

## Storage

Store the CD22 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).