


# CD28 Antibody (RQ7067)


Catalog No.	Formulation	Size
RQ7067	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P10747
Applications	Western Blot : 0.5-1ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This CD28 antibody is available for research use only.



Flow cytometry testing of human HEL cells with CD28 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= CD58 antibody.



Western blot testing of human 1) Jurkat and 2) HUT-78 cell lysate with CD28 antibody. Expected molecular weight depending on level of glycosylation: ~25-44 kDa (monomer), ~50-90kDa (dimer).

## **Description**

The protein encoded by this gene is essential for T-cell proliferation and survival, cytokine production, and T-helper type-2 development. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene.

#### **Application Notes**

Optimal dilution of the CD28 antibody should be determined by the researcher.

### **Immunogen**

Recombinant human protein (amino acids Q15-D214) was used as the immunogen for the CD28 antibody.

#### **Storage**

After reconstitution, the CD28 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.