


CD30 Antibody [clone CDLA30-1] (V7036)

Catalog No.	Formulation	Size
V7036-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7036-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7036SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7036IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	CDLA30-1
Purity	Protein G affinity chromatography
UniProt	P28908
Localization	Cell surface
Applications	Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT (1) Prediluted IHC Only Format: incubate for 30 min at RT (2)
Limitations	This CD30 antibody is available for research use only.



IHC testing of FFPE human Hodgkin's lymphoma and CD30 antibody (clone CDLA30-1).

Description

CD30 is the receptor for TNFSF8/CD30L. May play a role in the regulation of cellular growth and transformation of activated lymphoblasts. Regulates gene expression through activation of NF-kappa-B. [UniProt]

Application Notes

Optimal dilution of the CD30 antibody should be determined by the researcher.

- 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min.
- 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant human protein was used as the immunogen for the CD30 antibody.

Storage

Store the CD30 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).