


## CD30 Antibody (V3276)

| Catalog No.    | Formulation  | Size   |
|----------------|--|--------|
| V3276-100UG    | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V3276-20UG     | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug  |
| V3276SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free                             | 100 ug |

### Bulk quote request

|                           |   |
|---------------------------|---|
| <b>Availability</b>       | 1-3 business days   |
| <b>Species Reactivity</b> | Human   |
| <b>Format</b>             | Purified  |
| <b>Host</b>               | Rabbit  |
| <b>Clonality</b>          | Polyclonal (rabbit origin)                                |
| <b>Isotype</b>            | Rabbit IgG  |
| <b>Purity</b>             | Protein A affinity chromatography                         |
| <b>UniProt</b>            | P28908  |
| <b>Localization</b>       | Cell surface, cytoplasmic                                 |
| <b>Applications</b>       | Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT |
| <b>Limitations</b>        | This CD30 antibody is available for research use only.    |



IHC testing of FFPE human Hodgkin's lymphoma stained with CD30 antibody. Required HIER: steam sections in pH9, 1mM EDTA for 10-20 min.

## Description

CD30 is the receptor for TNFSF8/CD30L. May play a role in the regulation of cellular growth and transformation of activated lymphoblasts. Regulates gene expression through activation of NF-kappa-B. [UniProt]

## **Application Notes**

Optimal dilution of the CD30 antibody should be determined by the researcher.

## **Immunogen**

A partial human protein was used as the immunogen for the CD30 antibody.

## **Storage**

Store the CD30 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).