
CD31 Antibody PECAM1/3525 / Mouse Monoclonal [clone PECAM1/3525] (V8410)


Catalog No.	Formulation	Size
V8410-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8410-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8410SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG
Clone Name	PECAM1/3525
Purity	Protein G affinity chromatography
UniProt	P16284
Localization	Cell surface and cytoplasm of endothelial cells
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
Limitations	This CD31 Antibody PECAM1/3525 is available for research use only.




CD31 Antibody PECAM1/3525 Immunohistochemistry Analysis. IHC staining of FFPE human tonsil with CD31 antibody (clone PECAM1/3525). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human ovary with CD31 antibody (clone PECAM1/3525). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using CD31 antibody (clone PECAM1/3525). These results demonstrate the foremost specificity of the PECAM1/3525 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free CD31 antibody (clone PECAM1/3525) as confirmation of integrity and purity.

Description

CD31 Antibody PECAM1/3525 detects CD31 (PECAM-1), a transmembrane glycoprotein member of the immunoglobulin supergene family of adhesion molecules. CD31 is expressed by stem cells of the hematopoietic system and is primarily used to identify and concentrate these cells for experimental studies as well as for bone marrow transplantation. Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 MAb reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence.

For a Protein Microarray Validated monoclonal antibody with CD31 as the primary designation, see our CD31 Antibody PECAM1/3527 / PECAM-1 Monoclonal Antibody.

Application Notes

Optimal dilution of the CD31 Antibody PECAM1/3525 should be determined by the researcher.

Immunogen

A portion of amino acids 625-738 from the human protein was used as the immunogen for the CD31 antibody.

Storage

Store the CD31 Antibody PECAM1/3525 at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Alternate Names

PECAM-1 antibody, PECAM1 antibody, CD31 antibody, Platelet Endothelial Cell Adhesion Molecule 1 antibody, EndoCAM antibody, PECA1 antibody, GPIIA antibody

Copyright © NSJ Bioreagents. All rights reserved.