


# CD36 Antibody [clone 1E8.] (V3004)

Catalog No.	Formulation	Size
V3004-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3004-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3004SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Citations (5)	)
---------------	---

# **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	1E8.
Purity	Protein G affinity chromatography
UniProt	P16671
Localization	Cell surface
Applications	Functional Studies (order BSA/sodium Azide-free Format) : Flow Cytometry : 0.5-1ug/10^6 cells Immunofluorescence : 0.5-1ug/ml
Limitations	This CD36 antibody is available for research use only.



SDS-PAGE Analysis of Purified, BSA-Free CD36 Antibody (clone 1E8). Confirmation of Integrity and Purity of the Antibody.

Recognizes a protein of 80kDa-90kDa, identified as CD36. It is expressed on platelets, monocytes and macrophages, microvascular endothelial cells, erythrocyte precursors, mammary epithelial cells, and some macrophage derived dendritic cells. CD36 acts as a receptor for thrombospondin (TSP), collagen types I, IV and V, P. falciparum malaria-infected erythrocytes, and sickle erythrocytes. It also functions as a scavenger receptor, mediating macrophage uptake of oxidized low-density lipoprotein (LDL) and recognition of apoptotic polymorphonuclear leukocytes (PMN). CD36 plays a role in platelet aggregation, macrophage foam cell development, inflammation, and the tissue ischemia observed in sickle cell disease and cerebral malaria. Note that 1-4% of Japanese and East Asia population lack CD36.

## **Application Notes**

Optimal dilution of the CD36 antibody should be determined by the researcher.

### **Immunogen**

Human CD36 from platelets was used as the immunogen for the CD36 antibody.

#### **Storage**

Store the CD36 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).