


CD3e Antibody [clone C3e/2478] (V3751)


Catalog No.	Formulation	Size
V3751-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3751-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3751SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	C3e/2478
Purity	Protein G affinity chromatography
UniProt	P07766
Localization	Cell surface and cytoplasmic
Applications	Flow Cytometry : 1-2ug/10 ⁶ cells ELISA : 2-4ug/ml (order BSA/azide-free format) Western Blot : 1-2ug/ml
Limitations	This CD3e antibody is available for research use only.




Flow cytometry testing of human Jurkat cells with CD3e antibody (clone C3e/2478); Red=isotype control, Blue= CD3e antibody.




Western blot testing of human Jurkat cells with CD3e antibody (clone C3e/2478).
Expected molecular weight ~23 kDa.

Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using CD3e antibody (clone C3e/2478). These results demonstrate the foremost specificity of the C3e/2478 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free CD3e antibody (clone C3e/2478) as confirmation of integrity and purity.

Description

This antibody recognizes the epsilon chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intracytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms.

Application Notes

Optimal dilution of the CD3e antibody should be determined by the researcher.

Immunogen

A portion of amino acids 23-119 were used as the immunogen for the CD3e antibody.

Storage

Store the CD3e antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).

