


# CD4 Antibody [clone RIV7] (V8208)


Catalog No.	Formulation	Size
V8208-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8208-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8208SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a, kappa
Clone Name	RIV7
Purity	Protein G affinity chromatography
UniProt	P06332
Localization	Cell surface
Applications	Functional Studies (order BSA/azide-free Format) : Flow Cytometry : 1-2ug/10^6 cells in 0.1ml Immunofluorescence : 1-2ug/ml
Limitations	This CD4 antibody is available for research use only.



Flow cytometry staining of lymphocyte gated human PBM cells with CD4 antibody; Gray=isotype control, Red= CD4 antibody.



SDS-PAGE analysis of purified, BSA-free CD4 antibody (clone RIV7) as confirmation of integrity and purity.

## **Description**

Recognizes a protein of 55kDa, identified as CD4. It is a membrane glycoprotein of T lymphocytes that interacts with major histocompatibility complex class II antigens and is also a receptor for the human immunodeficiency virus. This protein is expressed not only in T lymphocytes, but also in B cells, macrophages, and granulocytes. It is also expressed in specific regions of the brain. The protein functions to initiate or augment the early phase of T-cell activation, and may function as an important mediator of indirect neuronal damage in infectious and immune-mediated diseases of the central nervous system. Multiple alternatively spliced transcript variants encoding different isoforms have been identified.

## **Application Notes**

Optimal dilution of the CD4 antibody should be determined by the researcher.

## **Immunogen**

Human peripheral lymphocytes were used as the immunogen for the CD4 antibody.

## **Storage**

Store the CD4 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).