


CD44 Antibody / HCAM [clone rHCAM/6449] (V8829)

Catalog No.	Formulation	Size
V8829-100UG	0.2~mg/ml in 1X PBS with $0.1~mg/ml$ BSA (US sourced), $0.05%$ sodium azide	100 ug
V8829-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V8829SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant MOUSE MONOCLONAL

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG2a, kappa
Clone Name	rHCAM/6449
Purity	Protein A/G affinity
UniProt	P16070
Localization	Cell Surface
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This CD44 antibody is available for research use only.



IHC staining of FFPE human esophagus tissue with CD44 antibody (clone rHCAM/6449). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human breast tissue with CD44 antibody (clone rHCAM/6449). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Negative control: IHC staining of FFPE human brain tissue using CD44 antibody (clone rHCAM/6449) at 2ug/ml in PBS for 30min RT. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Description

The CD44 family of glycoproteins exists in a number of variant isoforms, the most common being the standard 85-95kDa or hematopoietic variant (CD44s). Higher molecular weight isoforms are described in epithelial cells (CD44v), which are believed to function in intercellular adhesion and stromal binding.

Application Notes

Optimal dilution of the CD44 antibody should be determined by the researcher.

Immunogen

Recombinant full-length human CD44 protein was used as the immunogen for the CD44 antibody.

Storage

Aliquot the CD44 antibody and store frozen at -200C or colder. Avoid repeated freeze-thaw cycles.