


CD44v4/5 Antibody [clone 3D2] (V8250)


Catalog No.	Formulation	Size
V8250-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8250-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8250SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	3D2
Purity	Protein G affinity chromatography
UniProt	P16070
Applications	Flow Cytometry : 1-2ug/10 ⁶ cells in 0.1ml Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This CD44v4/5 antibody is available for research use only.



IHC staining of FFPE human cervix with CD44v4/5 antibody (clone 3D2). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human cervix with CD44v4/5 antibody (clone 3D2). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free CD44v4/5 antibody (clone 3D2) as confirmation of integrity and purity.

Description

This antibody recognizes an epitope encoded by exon v9 on the variant portion of human CD44. The CD44 molecule belongs to a family of cellular adhesion molecules found on a wide range of normal and malignant cells in epithelial, mesothelial and hemopoiesis tissues. CD44 is a single gene with 20 exons, of which 10 are normally expressed to encode the basic CD44 (H-CAM) molecule. The additional 10 exons (v1 to v10) are only expressed by alternative splicing of the nuclear RNA. The expression of specific cell adhesion molecule CD44 splice variants has been reported to be associated with metastasis in certain human malignancies.

Application Notes

Optimal dilution of the CD44v4/5 antibody should be determined by the researcher.

Immunogen

A recombinant human full-length CD44v3-10 protein was used as the immunogen for this CD44v4/5 antibody.

Storage

Store the CD44v4/5 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).