


# **CD44v6 Antibody [clone CD44V6/2496] (V3775)**


Catalog No.	Formulation	Size
V3775-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3775-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3775SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	CD44V6/2496
Purity	Protein G affinity chromatography
UniProt	P16070
Localization	Cell surface, cytoplasmic
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Limitations	This CD44v6 antibody is available for research use only.



IHC testing of FFPE human esophagus tissue with CD44v6 antibody (clone CD44V6/2496). Required HIER: steam sections in pH 9 10mM Tris with 1mM EDTA buffer for 10-20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free CD44v6 antibody (clone CD44V6/2496) as confirmation of integrity and purity.

### **Description**

This antibody recognizes an epitope encoded by exon v6 on the variant portion of human CD44. The CD44 molecule belongs to a family of cellular adhesion molecules found on a wide range of normal and malignant cells in epithelial, mesothelial and hemopoiesis tissues. CD44 is a single gene with 20 exons, of which 10 are normally expressed to encode the basic CD44 (H-CAM) molecule. The additional 10 exons (v1 to v10) are only expressed by alternative splicing of the nuclear RNA. The expression of specific cell adhesion molecule CD44 splice variants has been reported to be associated with metastasis in certain human malignancies.

## **Application Notes**

Titering of the CD44v6 antibody may be required for optimal performance.

#### **Immunogen**

An amino acid sequence from the variant 6 domain of CD44 was used as the immunogen for the CD44v6 antibody.

#### **Storage**

Store the CD44v6 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).