
CD45RA Antibody SPM504 / Mouse Monoclonal [clone SPM504] (V2413)


Catalog No.	Formulation	Size
V2413-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2413-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2413SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2413IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Bulk quote request

Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	SPM504
Purity	Protein G affinity chromatography
UniProt	P08575
Localization	Cell surface
Applications	Western blot: 1-2ug/ml Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT (1) (2)
Limitations	This CD45RA Antibody SPM504 is available for research use only.



CD45RA Antibody SPM504 WB. Western blot testing of human spleen lysate with CD45RA antibody (clone SPM504). Expected molecular weight: 147-220 kDa depending on glycosylation level.



CD45RA Antibody SPM504 Immunohistochemistry. IHC test of FFPE human tonsil probed with CD45RA antibody (clone SPM504).

Description

CD45RA Antibody SPM504 detects CD45RA, an isoform of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C, respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD4+ T-cells, 50% of peripheral CD8+ T-cells, B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory T-cells. CD45RA and CD45RO define complementary, predominantly non-overlapping populations of resting peripheral T-cells. This MAb is useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especially be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphomas (CD45RA +ve).

CD45RA is an alternatively spliced form of PTPRC widely studied in lymphocyte differentiation and immune cell phenotyping; learn more about the target and related reagents on our CD45RA Antibody page.

Application Notes

Due to variation in protocol and secondary antibody used, the CD45RA Antibody SPM504 may need to be titered for optimal performance.

1. FFPE staining requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Stimulated human leukocytes were used as immunogen for this CD45RA antibody.

Storage

The CD45RA antibody (with azide) can be stored at 2-8oC. The azide-free format should be aliquoted and stored at -20oC or colder.

References (2)

Copyright © NSJ Bioreagents. All rights reserved.