


## CD47 Antibody / IAP [clone IAP/964] (V3014)


Catalog No.	Formulation	Size
V3014-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3014-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3014SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Bulk quote request


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	IAP/964
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	Q08722
<b>Localization</b>	Cell surface, cytoplasmic
<b>Applications</b>	Immunofluorescence : 1-2ug/ml Flow Cytometry : 1-2ug/million cells
<b>Limitations</b>	This CD47 antibody is available for research use only.




Immunofluorescent staining of live human Jurkat cells with CD47 antibody (green, clone IAP/964) and Reddot nuclear stain (red).



Immunofluorescent staining of PFA-fixed human MCF7 cells with CD47 antibody (green, clone IAP/964) and Reddot nuclear stain (red).



Flow cytometry staining of live human Jukat cells with CD47 antibody (clone IAP/964); Red=isotype control, Blue= CD47 antibody.



SDS-PAGE analysis of purified, BSA-free CD47 antibody (clone IAP/964) as confirmation of integrity and purity.

## Description

This antibody reacts with Ig domain of CD47 protein. CD47, originally named integrin-associated protein (IAP), is a 50kDa protein containing five membrane-spanning sequences and a short cytoplasmic tail. CD47 plays a role in both cell adhesion by acting as an adhesion receptor for THBS1 on platelets, and in the modulation of integrins. It is important in memory formation and synaptic plasticity in the hippocampus. CD47 may play a role in membrane transport and/or integrin dependent signal transduction.

## Application Notes

Optimal dilution of the CD47 antibody should be determined by the researcher.

## Immunogen

Recombinant human protein was used as the immunogen for the CD47 antibody.

## Storage

Store the CD47 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).

