


## CD57 Antibody / HNK-1 [clone NK1/7565] (V4220)

Catalog No.	Formulation	Size
V4220-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4220-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4220SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


### Bulk quote request

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2, lambda
<b>Clone Name</b>	NK1/7565
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	Q9P2W7
<b>Localization</b>	Cell surface, cytoplasmic
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
<b>Limitations</b>	This CD57 antibody is available for research use only.



IHC staining of FFPE human stomach tissue with CD57 antibody (clone NK1/7565).  
 HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

SDS-PAGE analysis of purified, BSA-free CD57 antibody (clone NK1/7565) as confirmation of integrity and purity.



## Description

Anti-CD57 marks a subset of lymphocytes known as natural killer (NK) cells. Follicular center cell lymphomas often contain many NK cells within the neoplastic follicles. Anti-CD57 also stains neuroendocrine cells and their derived tumors, including carcinoid tumor and medulloblastoma. Anti-CD57 can also be useful in separating type B3 thymoma from thymic carcinoma when combined with a panel that includes antibodies against GLUT1, CD5, and CEA.

## Application Notes

Optimal dilution of the CD57 antibody should be determined by the researcher.

## Immunogen

A recombinant partial protein (within amino acids 1-200) from the human protein was used as the immunogen for the CD57 antibody.

## Storage

Aliquot the CD57 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.