


CD61 Antibody / ITGB3 / Integrin beta 3 (RQ7183)


Catalog No.	Formulation	Size
RQ7183	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P05106
Localization	Cell surface, cytoplasmic
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This CD61 antibody is available for research use only.




IHC staining of FFPE human lung cancer tissue with CD61 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human liver cancer tissue with CD61 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE rat lung tissue with CD61 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) human U-87 MG, 2) human HEL and 3) rat spleen lysate with CD61 antibody. Expected molecular weight: 87-110 kDa depending on glycosylation level.



Flow cytometry testing of human HEL cells with CD61 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= CD61 antibody.

Description

ITGB3 (INTEGRIN, BETA-3), also called GP3A, GPIIIa, CD61, is a protein that in humans is encoded by the ITGB3 gene. It is a cluster of differentiation found on thrombocytes. The 3-prime exon is larger than 1,700 nucleotides and contains the 3-prime untranslated region. The ITGB3 complex belongs to the integrin class of cell adhesion molecule receptors that share a common heterodimeric structure with alpha and beta subunits. Additionally, the ITGB3 complex mediates platelet aggregation by acting as a receptor for fibrinogen. Although the ITGB3 is expressed on the cell surface at normal levels and is capable of function following extracellular stimulation, it could not be activated via the 'inside-out' signaling pathways.

Application Notes

Optimal dilution of the CD61 antibody should be determined by the researcher.

Immunogen

Recombinant human protein (amino acids N29-Q523) was used as the immunogen for the CD61 antibody.

Storage

After reconstitution, the CD61 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.