


## CD63 Antibody / LAMP-3 [clone LAMP3/7369] (V4263)


Catalog No.	Formulation	Size
V4263-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4263-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4263SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Bulk quote request

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2, kappa
<b>Clone Name</b>	LAMP3/7369
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	P08962
<b>Localization</b>	Cell surface, Cytoplasm
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
<b>Limitations</b>	This CD63 antibody is available for research use only.



IHC staining of FFPE human melanoma tissue with CD63 antibody (clone LAMP3/7369).  
 HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free CD63 antibody (clone LAMP3/7369) as confirmation of integrity and purity.

## Description

CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression.

## Application Notes

Optimal dilution of the CD63 antibody should be determined by the researcher.

## Immunogen

A recombinant partial protein sequence (within amino acids 100-200) from the human protein was used as the immunogen for the CD63 antibody.

## Storage

Aliquot the CD63 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.