


CD64 Antibody / FCGR1A (RQ7170)


Catalog No.	Formulation	Size
RQ7170	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P12314
Applications	Western Blot : 0.5-1ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This CD64 antibody is available for research use only.



Flow cytometry testing of human SiHa cells with CD64 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= CD64 antibody.



Western blot testing of 1) human ThP-1, 2) rat spleen, 3) rat thymus, 4) rat PC-3, 5) mouse EL-4, 6) mouse spleen, 7) mouse thymus, 8) mouse RAW264.7 and 9) mouse ANA-1 cell lysate with CD64 antibody. Predicted molecular weight: 39-75 kDa depending on glycosylation level.

Description

High affinity immunoglobulin gamma Fc receptor I is a protein that in humans is encoded by the FCGR1A gene. It is mapped to 1q21.2. This gene encodes a protein that plays an important role in the immune response. This protein is a high-affinity Fc-gamma receptor. The gene is one of three related gene family members located on chromosome 1.

Application Notes

Optimal dilution of the CD64 antibody should be determined by the researcher.

Immunogen

Recombinant human protein (amino acids A191-D254) was used as the immunogen for the CD64 antibody.

Storage

After reconstitution, the CD64 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.