


CD64 Antibody [clone FCGR1A/7498] (V4723)


Catalog No.	Formulation	Size
V4723-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4723-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4723SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG
Clone Name	FCGR1A/7498
Purity	Protein A/G affinity
UniProt	P12314
Localization	Cell Membrane
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This CD64 antibody is available for research use only.



IHC staining of FFPE human tonsil tissue using CD64 antibody (clone FCGR1A/7498) at 2ug/ml. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free CD64 antibody (clone FCGR1A/7498) as confirmation of integrity and purity.

Description

CD64 is a surface glycoprotein with high affinity for monomeric IgG, is expressed constitutively on monocytes and macrophages, and can be induced in neutrophils subsequent to IFN-gamma stimulation.

Application Notes

Optimal dilution of the CD64 antibody should be determined by the researcher.

Immunogen

Recombinant human FCGR1A protein was used as the immunogen for the CD64 antibody.

Storage

Aliquot the CD64 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.