


# CD68 Antibody [clone CD68/G2] (V3032)

Catalog No.	Formulation	Size
V3032-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3032-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3032SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3032IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

# Citations (1)

### **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	CD68/G2
Purity	Protein G affinity chromatography
UniProt	P34810
Localization	Cytoplasmic, membrane
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Limitations	This CD68 antibody is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human Histiocytoma stained with CD68 antibody (CD68/G2).

#### **Description**

This antibody recognizes a glycoprotein of 110kDa, which is identified as CD68. It is important for identifying macrophages in tissue sections. It stains macrophages in a wide variety of human tissues, including Kupffer cells and macrophages in the red pulp of the spleen, in lamina propria of the gut, in lung alveoli, and in bone marrow. It reacts with myeloid precursors and peripheral blood granulocytes. It also reacts with plasmacytoid T cells, which are supposed to be of monocyte/macrophage origin. It shows strong granular cytoplasmic staining of chronic and acute myeloid leukemia and also reacts with rare cases of true histiocytic neoplasia. Lymphomas are negative or show few granules.

#### **Application Notes**

Optimal dilution of the CD68 antibody should be determined by the researcher.

- 1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.
- 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

### **Immunogen**

Recombinant human protein was used as the immunogen for the CD68 antibody.

#### **Storage**

Store the CD68 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).