


# **CD7 Antibody [clone 124-1D1] (V2964)**

Catalog No.	Formulation	Size
V2964-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2964-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2964SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	124-1D1
Purity	Protein G affinity chromatography
UniProt	P09564
Localization	Cell surface
Applications	Flow Cytometry : 0.5-1ug/10^6 cells Immunofluorescence : 0.5-1ug/ml
Limitations	This CD7 antibody is available for research use only.



SDS-PAGE Analysis of Purified, BSA-Free CD7 Antibody (clone 124-1D1). Confirmation of Integrity and Purity of the Antibody.

## **Description**

gene superfamily. Its N-terminal amino acids 1-107 are highly homologous to Ig kappa-L chains whereas the carboxyl-terminal region of the extracellular domain is proline-rich and has been postulated to form a stalk from which the Ig domain projects. CD7 is expressed on the majority of immature and mature T-lymphocytes, and T cell leukemia. It is also found on natural killer cells, a small subpopulation of normal B cells and on malignant B cells. Cross-linking surface CD7 positively modulates T cell and NK cell activity as measured by calcium fluxes, expression of adhesion molecules, cytokine secretion and proliferation. CD7 associates directly with phosphoinositol 3'-kinase. CD7 ligation induces production of D-3 phosphoinositides and tyrosine phosphorylation.

#### **Application Notes**

Optimal dilution of the CD7 antibody should be determined by the researcher.

### **Immunogen**

Human leukocytes were used as the immunogen for the CD7 antibody.

#### **Storage**

Store the CD7 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).