


# **CD7 Antibody [clone CD7/8118R] (V4559)**

Catalog No.	Formulation	Size
V4559-100UG	0.2~mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4559-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4559SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


### Recombinant RABBIT MONOCLONAL

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	CD7/8118R
Purity	Protein A/G affinity
UniProt	P09564
Localization	Cell Surface
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Limitations	This CD7 antibody is available for research use only.



IHC staining of FFPE human tonsil tissue with CD7 antibody (clone CD7/8118R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human tonsil tissue with CD7 antibody (clone CD7/8118R). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

#### **Description**

Recognizes a protein of 40kDa, identified as CD7 (also known as gp40, Leu9). CD7 is a member of the immunoglobulin gene superfamily. Its N-terminal amino acids 1-107 are highly homologous to Ig kappa-L chains whereas the carboxylterminal region of the extracellular domain is proline-rich and has been postulated to form a stalk from which the Ig domain projects. CD7 is expressed on the majority of immature and mature T-lymphocytes, and T cell leukemia. It is also found on natural killer cells, a small subpopulation of normal B cells and on malignant B cells. Cross-linking surface CD7 positively modulates T cell and NK cell activity as measured by calcium fluxes, expression of adhesion molecules, cytokine secretion and proliferation. CD7 associates directly with phosphoinositol 3'-kinase. CD7 ligation induces production of D-3 phosphoinositides and tyrosine phosphorylation.

#### **Application Notes**

Optimal dilution of the CD7 antibody should be determined by the researcher.

#### **Immunogen**

A recombinant partial protein sequence (within amino acids 1-240) from the human protein was used as the immunogen for the CD7 antibody.

#### **Storage**

Aliquot the CD7 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.