


CD7 Antibody [clone HuLy-m2] (V2963)

Catalog No.	Formulation	Size
V2963-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2963-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2963SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Citations	(3)	
Ontationio	(\cup)	

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a, kappa
Clone Name	HuLy-m2
Purity	Protein G affinity chromatography
UniProt	P09564
Localization	Cell surface
Applications	Flow Cytometry : 1-2ug/10^6 cells Immunofluorescence : 1-3ug/ml
Limitations	This CD7 antibody is available for research use only.



SDS-PAGE analysis of purified, BSA-free CD7 antibody (clone HuLy-m2) as confirmation of integrity and purity.

Description

gene superfamily. Its N-terminal amino acids 1-107 are highly homologous to Ig kappa-L chains whereas the carboxyl-terminal region of the extracellular domain is proline-rich and has been postulated to form a stalk from which the Ig domain projects. CD7 is expressed on the majority of immature and mature T-lymphocytes, and T cell leukemia. It is also found on natural killer cells, a small subpopulation of normal B cells and on malignant B cells. Cross-linking surface CD7 positively modulates T cell and NK cell activity as measured by calcium fluxes, expression of adhesion molecules, cytokine secretion and proliferation. CD7 associates directly with phosphoinositol 3'-kinase. CD7 ligation induces production of D-3 phosphoinositides and tyrosine phosphorylation.

Application Notes

Optimal dilution of the CD7 antibody should be determined by the researcher.

Immunogen

Human thymocytes were used as the immunogen for the CD7 antibody.

Storage

Store the CD7 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).