


CD73 Antibody (F47794)


Catalog No.	Formulation	Size
F47794-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F47794-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human, Mouse
Format	Purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Purified
UniProt	P21589
Applications	Western Blot : 1:1000-2000 IHC (Paraffin) : 1:10-1:50
Limitations	This CD73 antibody is available for research use only.



Western blot analysis of lysate from (1) U266, (2) HepG2 cell line, (3) mouse brain tissue lysate using CD73 antibody at 1:2000. Predicted molecular weight: 65-70 kDa.



Western blot analysis of NT5E/ CD73 antibody and 293 cell lysate (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (2) with the NT5E gene.



IHC analysis of FFPE human placenta tissue stained with CD73 antibody

Description

Ecto-5-prime-nucleotidase/5-prime-ribonucleotide phosphohydrolase/NT5E/CD73 catalyzes the conversion at neutral pH of purine 5-prime mononucleotides to nucleosides, the preferred substrate being AMP. The enzyme consists of a dimer of 2 identical 70 kD subunits bound externally to the plasma membrane by a glycosyl phosphatidyl inositol linkage. The enzyme is used as a marker of lymphocyte differentiation. Consequently, a deficiency of NT5E/CD73 occurs in a variety of immunodeficiency diseases. Other forms of 5-prime nucleotidase exist in the cytoplasm and lysosomes and can be distinguished from ecto-NT5 by their substrate affinities, requirement for divalent magnesium ion, activation by ATP, and inhibition by inorganic phosphate. It is not known whether the different enzymes are coded by different genes or result from different posttranslational modifications of a single coding sequence.

Application Notes

Titration of the CD73 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 520-550 from the human protein was used as the immunogen for this CD73 antibody.

Storage

Aliquot the CD73 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.