


## CD74 Antibody / CLIP [clone CLIP/7194] (V5103)


Catalog No.	Formulation	Size
V5103-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5103-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5103SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	CLIP/7194
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	P04233
<b>Localization</b>	Cell surface, Paranuclear globular
<b>Applications</b>	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
<b>Limitations</b>	This CD74 antibody is available for research use only.



IHC staining of FFPE human spleen tissue with CD74 antibody (clone CLIP/7194) at 2ug/ml. Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Western blot testing of human Raji cell lysate with CD74 antibody (clone CLIP/7194).  
Expected molecular weight: 33-43 kDa.

## Description

It recognizes a protein of ~35kDa, identified as CD74 (Workshop IV). CD74 is a type II transmembrane protein which binds to the peptide binding groove of newly synthesized MHC class II alpha/beta heterodimers and prevents their premature association with endogenous polypeptides. CD74 is expressed primarily by antigen presenting cells, such as B-lymphocytes (from before the pre-B cell stage to before the plasma cell stage), macrophages, and monocytes, and many epithelial cells. Anti-CD74 stains predominantly germinal center lymphocytes and B-cell lymphomas, but rarely T-cell lymphomas. Anti-CD74 has been shown to be useful in differentiating atypical fibroxanthoma (-) from malignant fibrous histiocytoma (+).

## Application Notes

Optimal dilution of the CD74 antibody should be determined by the researcher.

## Immunogen

A recombinant fragment of human CD74 protein was used as the immunogen for the CD74 antibody.

## Storage

Aliquot the CD74 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.