


CD74 Antibody (RQ7512)


Catalog No.	Formulation	Size
RQ7512	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P04233
Localization	Cell surface and paranuclear globular
Applications	Western Blot: 0.5-1ug/ml Immunohistochemistry (FFPE): 2-5ug/ml Immunofluorescence: 5ug/ml Flow Cytometry: 1-3ug/million cells Direct ELISA: 0.1-0.5ug/ml
Limitations	This CD74 antibody is available for research use only.




IHC staining of FFPE human spleen tissue with CD74 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human tonsil tissue with CD74 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Immunofluorescent staining of FFPE human spleen tissue with CD74 antibody (red) and DAPI nuclear stain (blue). HIER: steam section in pH8 EDTA buffer for 20 min.



Western blot testing of human Daudi cell lysate with CD74 antibody. Expected molecular weight: 33-43 kDa.



Flow cytometry testing of human Daudi cells with CD74 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= CD74 antibody.

Description

HLA class II histocompatibility antigen gamma chain also known as HLA-DR antigens-associated invariant chain or CD74 (Cluster of Differentiation 74), is a protein that in humans is encoded by the CD74 gene. The protein encoded by this gene associates with class II major histocompatibility complex (MHC) and is an important chaperone that regulates antigen presentation for immune response. It also serves as cell surface receptor for the cytokine macrophage migration inhibitory factor (MIF) which, when bound to the encoded protein, initiates survival pathways and cell proliferation. This protein also interacts with amyloid precursor protein (APP) and suppresses the production of amyloid beta (Abeta). Multiple alternatively spliced transcript variants encoding different isoforms have been identified.

Application Notes

Optimal dilution of the CD74 antibody should be determined by the researcher.

Immunogen

E. coli-derived recombinant human protein (amino acids Q73-M296) was used as the immunogen for the CD74 antibody.

Storage

After reconstitution, the CD74 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.