
CD79a Antibody [clone IGA/1406] (V8448)


Catalog No.	Formulation	Size
V8448-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8448-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8448SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG3, kappa
Clone Name	IGA/1406
Purity	Protein G affinity chromatography
UniProt	P11912
Localization	Cell surface, cytoplasmic
Applications	Western blot: 1-2ug/ml
Limitations	This CD79a antibody is available for research use only.



Western blot testing of human Ramos cell lysate with CD79a antibody (clone IGA/1406). Expected molecular weight: 25-47 kDa depending on glycosylation level.



SDS-PAGE analysis of purified, BSA-free CD79a antibody (clone IGA/1406) as confirmation of integrity and purity.

Description

A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.

Application Notes

Optimal dilution of the CD79a antibody should be determined by the researcher.

Immunogen

Amino acids GTYQDVGSLNIADVQ were used as the immunogen for the CD79a antibody.

Storage

Store the CD79a antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Ordering: Phone: 858.663.9055 | Fax: 1.267.821.0800 | Email: info@nsjbio.com
Copyright © NSJ Bioreagents. All rights reserved.