


# CD80 Antibody (R30433)

Catalog No.	Formulation	Size
R30433	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

### **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P33681
Localization	Cytoplasmic, membranous
Applications	Western Blot : 0.5-1ug/ml
Limitations	This CD80 antibody is available for research use only.



Western blot testing of human 1) Raji, 2) Daudi and 3) Ramos cell lysate with CD80 antibody. Expected molecular weight: 34-75 kDa depending on the level of glycosylation.

# **Description**

Cluster of Differentiation 80 (also CD80 and B7-1) is a protein found on activated B cells and monocytes that provides a costimulatory signal necessary for T cell activation and survival. It is the ligand for two different proteins on the T cell surface: CD28 (for autoregulation and intercellular association) and CTLA-4 (for attenuation of regulation and cellular disassociation). CD80 works in tandem with CD86 to prime T cells. The CD80 genes encode B7-1 which are structurally similar members of the immunoglobulin superfamily expressed on a variety of hematopoietic cell types. Reeves et al.(1997) stated that B7-1 and B7-2 provide a costimulatory signal to T cells by interacting with CD28 and CTLA4.

# **Application Notes**

The stated application concentrations are suggested starting amounts. Titration of the CD80 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

#### **Immunogen**

An amino acid sequence from the N-terminus of human CD80 (EELAQTRIYWQKEKK) was used as the immunogen for this CD80 antibody.

#### **Storage**

After reconstitution, the CD80 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.