

CD99 Antibody [clone SPM586] (V2707)

Catalog No.	Formulation	Size
V2707-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2707-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2707SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2707IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgM, kappa
Clone Name	SPM586
Purity	PEG precipitation
UniProt	P14209
Localization	Cell surface
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Limitations	This CD99 antibody is available for research use only.

IHC: Formalin-fixed, paraffin-embedded human Ewing's sarcoma stained with CD99 antibody (SPM586).

Description

Recognizes a sialoglycoprotein of 27-32kDa, identified as CD99, or MIC2 gene product, or E2 antigen. This antigen is expressed on the cell membrane of some lymphocytes, cortical thymocytes, and granulosa cells of the ovary. Most pancreatic islet cells, Sertoli cells of the testis, and some endothelial cells express this antigen. Mature granulocytes express very little or no CD99. MIC2 is strongly expressed on Ewing s sarcoma cells and primitive peripheral neuroectodermal tumors. This mAb shows a very similar reactivity to other CD99 mAbs (e.g. O13, 12E7, or HBA-71) and

is excellent for immunohistochemical staining of formalin-fixed, paraffin-embedded tissues.

Application Notes

Optimal dilution of the CD99 antibody should be determined by the researcher.

- 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min
- 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Purified E-rosette forming cells from human peripheral blood lymphocytes were used as the immunogen for the CD99 antibody.

Storage

Store the CD99 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).