
Cdc20 Antibody [clone AR12] (V2084)

Catalog No.	Formulation	Size
V2084-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2084-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2084SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2084IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Citations (3)

Bulk quote request


Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	AR12
Purity	Protein G purified antibody
Buffer	1X PBS, pH 7.4
Gene ID	991
Localization	Cytoplasmic
Applications	Flow cytometry: 1-2ug/10^6 cells Immunofluorescence: 1-2ug/ml Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Limitations	This **Cdc20 antibody** is available for research use only.



IHC staining of FFPE human tonsil with Cdc20 antibody (clone AR12). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human gastric carcinoma with Cdc20 antibody (clone AR12). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Cdc20 antibody (clone AR12) as confirmation of integrity and purity.

Description

Cyclins, regulatory subunits which associate with kinases, control many of the important steps in cell cycle progression. The Cdc2 kinase (p34Cdc2) exhibits kinase activity in vitro and exists in a complex with both cyclin B and a protein homologous to p13SUC1. Cdc2 kinase is the active subunit of the M phase promoting factor (MPF) and the M phase-specific Histone H1 kinase. The p34Cdc2/cyclin B complex is required for the G2 to M transition. An additional cell cycle-dependent protein kinase, termed p55cdc, exhibits a high degree of homology with the S. cerevisiae proteins Cdc20 and Cdc4. The p55cdc transcript is readily detectable in a variety of cultured cell lines in growth phase, but disappears when cell growth is chemically arrested.

Application Notes

The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the Cdc20 antibody to be titered up or down for optimal performance.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Denatured recombinant human protein was used as the immunogen for this Cdc20 antibody.

Storage

Store the Cdc20 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Alternate Names

CDC20A, p55CDC, P55CDC-LSB, Cdc20 antibody

References (1)

Copyright © NSJ Bioreagents. All rights reserved.