
CDK4 Antibody [clone CDK4/8210R] (V4307)

Catalog No.	Formulation	Size
V4307-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4307-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4307SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant RABBIT MONOCLONAL

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	CDK4/8210R
Purity	Protein A/G affinity
UniProt	P11802
Localization	Nucleus, Cytoplasm
Applications	Immunohistochemistry (FFPE): 1-2ug/ml Western blot: 2-4ug/ml
Limitations	This CDK4 antibody is available for research use only.




CDK4 Antibody K-562 WB. Western blot analysis of Cyclin-dependent kinase 4 (CDK4) expression in K-562 cell lysate using CDK4 antibody clone CDK4/8210R. Lane 1: K-562 cell lysate. A band is detected at approximately 33-34 kDa, consistent with the predicted molecular weight of CDK4, a key regulator of G1 to S phase cell cycle progression. A faint lower-molecular-weight band is also observed below 20 kDa, which may represent proteolytic processing or non-specific reactivity.



CDK4 Antibody Human Tonsil Tissue IHC. Immunohistochemistry staining of FFPE human tonsil tissue with CDK4 antibody (clone CDK4/8210R). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free CDK4 antibody (CDK4/8210R) as confirmation of integrity and purity.



CDK4 Antibody Cell Line WB. Western blot analysis of Cyclin-dependent kinase 4 (CDK4) expression in cell lysates using CDK4 antibody. Lane 1: MCF7 cell lysate, Lane 2: HEK293T cell lysate. A band is detected at approximately 33â€“34 kDa, consistent with the predicted molecular weight of CDK4, a key regulator of G1 to S phase cell cycle progression. A lower-molecular-weight band is also observed at approximately 15â€“20 kDa, which may represent proteolytic processing or non-specific reactivity. The presence of CDK4 signal in both cell lines aligns with its known expression in proliferating cells.

Description

Cyclin-dependent kinase-4 (CDK4) is a protein-serine kinase involved in the cell cycle. It is essential for the G1- to S-phase transition during the cell cycle and its expression is primarily controlled at the transcriptional level. CCND1- CDK4 axis is not only critical in glial tumor cells but also in stromal-derived cells in the surrounding tumor microenvironment that are vital to sustain tumor outgrowth. CDK4 is highly expressed in highly differentiated and dedifferentiated liposarcomas, but rarely expressed in other benign liposarcomas and other sarcomas. CDK4 and MDM2 combined to differentiate between highly differentiated liposarcoma (+), dedifferentiated liposarcoma (+) and myxoid liposarcoma, pleomorphic liposarcoma, spindle lipoma, pleomorphic lipoma and other high-grade sarcomas.

This antibody is part of a broader antibody panel offered by NSJ Bioreagents.

Application Notes

Optimal dilution of the CDK4 antibody should be determined by the researcher.

Immunogen

A recombinant fragment corresponding to the N-terminal of human CDK4 protein was used as the immunogen for the CDK4 antibody.

Storage

Aliquot the CDK4 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

Copyright © NSJ Bioreagents. All rights reserved.