


Cdk4 Antibody (F52368)


Catalog No.	Formulation	Size
F52368-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F52368-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity
UniProt	P35426
Localization	Nuclear and cytoplasmic
Applications	IHC (Paraffin) : 1:25 Immunofluorescence : 1:25
Limitations	This Cdk4 antibody is available for research use only.



IHC testing of paraffin-embedded rat small intestine using Cdk4 antibody at 1:25.



IHC analysis of FFPE rat brain section using Cdk4 antibody at 1:25.



Fluorescent image of MCF-7 cells stained with Cdk4 antibody. Ab was diluted at 1:25 dilution. An Alexa Fluor 488-conjugated goat anti-rabbit IgG was used as the secondary Ab (green). Cytoplasmic actin was counterstained with Alexa Fluor 555 conjugated with Phalloidin (red).

Description

Ser/Thr-kinase component of cyclin D-CDK4 (DC) complexes that phosphorylate and inhibit members of the retinoblastoma (RB) protein family including RB1 and regulate the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complexes and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase. Hypophosphorylates RB1 in early G(1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals. Also phosphorylates SMAD3 in a cell-cycle-dependent manner and represses its transcriptional activity. Component of the ternary complex, cyclin D/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex (By similarity).

Application Notes

Titration of the Cdk4 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 272-303 from the rat protein was used as the immunogen for this Cdk4 antibody.

Storage

Aliquot the Cdk4 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.