


CDK9 Antibody (C-Terminal Region) (R36496)

| Catalog No. | Formulation | Size |
|--------------|---|--------|
| R36496-100UG | 0.5 mg/ml in 1X TBS, pH7.3, with 0.5% BSA (US sourced) and 0.02% sodium azide | 100 ug |

Bulk quote request

| | |
|-----------------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Mouse |
| Predicted Reactivity | Cow, Dog, Human, Pig, Rat |
| Format | Antigen affinity purified |
| Host | Goat |
| Clonality | Polyclonal (goat origin) |
| Isotype | Goat Ig |
| Purity | Antigen affinity |
| UniProt | Q99J95 |
| Applications | Western Blot : 0.3-1ug/ml ELISA (peptide) LOD : 1:8000 |
| Limitations | This CDK9 antibody is available for research use only. |



Western blot testing of mouse NIH3T3 cell lysate with CDK9 antibody at 2ug/ml.
Predicted molecular weight: ~43 kDa.

Description

Alternate name(s) for this target: Cyclin-dependent kinase 9; C-2k; CDC2L4; CTK1; PITALR; TAK; CDC2-related kinase, cell division cycle 2-like protein kinase 4; cell division protein kinase 9; serine/threonine protein kinase PITALRE; tat-associated kinase complex catalytic subunit

Application Notes

Optimal dilution of the CDK9 antibody should be determined by the researcher.

Immunogen

Amino acids KGSQITQQSTNQSRN were used as the immunogen for this CDK9 antibody.

Storage

Aliquot and store the CDK9 antibody at -20oC.