



## CDX2 Antibody [clone PCRP-CDX2-1A3] (V8501)

Catalog No.	Formulation	Size
V8501-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8501-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8501SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


### Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	PCRP-CDX2-1A3
Purity	Protein G affinity chromatography
UniProt	Q99626
Localization	Nuclear
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This CDX2 antibody is available for research use only.





IHC staining of FFPE human colon carcinoma with CDX2 antibody (clone PCRP-CDX2-1A3). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free CDX2 antibody (clone PCRP-CDX2-1A3) as confirmation of integrity and purity.

## Description

The intestine-specific transcription factors CDX1 and CDX2 are important for directing intestinal development, differentiation, proliferation and maintenance of the intestinal phenotype. CDX2 protein expression has been seen in GI carcinomas. Anti-CDX2 has been useful to establish GI origin of metastatic adenocarcinomas and carcinoids and is especially useful to distinguish metastatic colorectal adenocarcinoma from lung adenocarcinoma. However, mucinous carcinomas of the ovary also express CDX2 protein. It limits the usefulness of this marker in the distinction of metastatic colorectal adenocarcinoma from mucinous carcinoma of the ovary.

## Application Notes

Optimal dilution of the CDX2 antibody should be determined by the researcher.

## Immunogen

Recombinant human full-length CDX2 protein was used as the immunogen for the CDX2 antibody.

## Storage

Store the CDX2 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).