


## CETN1 Antibody / Centrin 1 [clone CETN1/9878] (V5656)

Catalog No.	Formulation	Size
V5656-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5656-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5656SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG
<b>Clone Name</b>	CETN1/9878
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	Q12798
<b>Localization</b>	Cell projection, Cytoplasm, Nucleus
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml
<b>Limitations</b>	This CETN1 antibody is available for research use only.



SDS-PAGE analysis of purified, BSA-free CETN1 antibody (clone CETN1/9878) as confirmation of integrity and purity.

### Description

EF-hand type Ca<sup>2+</sup>-binding proteins consists of several family members, including Centrin-1, Centrin-2 and Centrin-3. The Centrin proteins are ubiquitously expressed cytoskeletal components that show increased expression during cell

differentiation. Tissues where cilia are present, such as the retina and testis, express both Centrin-1 and -2, but Centrin-2 is also expressed in nondifferentiated, nonciliated retinal cells (retinoblastoma cells), liver, skeletal muscle, and cardiac muscle. In these tissues, Centrin associates with the centrosomes, mitotic spindle poles, and basal bodies. Knockdown studies reveal a requirement for Centrin in centriole duplication and organization of spindle pole morphology and the completion of cytokinesis. Centrin-3 plays a role in centrosome reproduction.

## **Application Notes**

Optimal dilution of the CETN1 antibody should be determined by the researcher.

## **Immunogen**

A portion of amino acids 1-172 from human CETN1 protein was used as the immunogen for the CETN1 antibody.

## **Storage**

Aliquot the CETN1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.