


CFTR Antibody / Cystic Fibrosis Transmembrane Regulator [clone CFTR/1342] (V7666)


Catalog No.	Formulation	Size
V7666-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7666-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7666SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human, Mouse
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a, kappa
Clone Name	CFTR/1342
Purity	Protein G affinity chromatography
UniProt	P13569
Localization	Cell surface, cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This CFTR antibody is available for research use only.



IHC testing of FFPE mouse kidney with CFTR antibody (clone CFTR/1342). Staining of FFPE tissue is enhanced by boiling tissue sections in 10mM Tris with 1mM EDTA, pH9 for 10-20 min followed by cooling at RT for 20 min.



IHC testing of FFPE human pancreas with CFTR antibody (clone CFTR/1342). Staining of FFPE tissue is enhanced by boiling tissue sections in 10mM Tris with 1mM EDTA, pH9 for 10-20 min followed by cooling at RT for 20 min.

Description

CFTR is composed of two membrane-spanning domains (MSD), two nucleotide-binding domains (NBD), and an R domain. It is structurally similar to multidrug resistance (Mdr1) protein and both are members of the superfamily of ATP-binding cassette (ABC) transporters, also known as traffic ATPases, which are implicated in the movement of various substrates. The CFTR protein is a small conductance adenosine 3',5'-cyclic monophosphate (cAMP)-activated chloride ion channel found in the apical membranes of epithelia within the pancreas, airway, intestine, bile duct, sweat gland, and male genital ducts. CFTR is a valuable marker of human pancreatic duct cell development and differentiation.

Application Notes

Optimal dilution of the CFTR antibody should be determined by the researcher.

Immunogen

Human recombinant protein was used as the immunogen for this CFTR antibody.

Storage

Store the CFTR antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).