


# Claudin 2 Antibody / CLDN2 (RQ6463)


Catalog No.	Formulation	Size
RQ6463	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

## **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P57739
Applications	Western Blot : 1-2ug/ml Immunofluorescence : 5ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This Claudin 2 antibody is available for research use only.



Immunofluorescent staining of FFPE human Caco-2 cells with Claudin 2 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Flow cytometry testing of human A549 cells with Claudin 2 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Claudin 2 antibody.



### **Description**

Claudin-2 is a protein that in humans is encoded by the CLDN2 gene. By genomic sequence analysis, this gene is mapped to chromosome Xq22.3-q23. This gene product belongs to the claudin protein family whose members have been identified as major integral membrane proteins localized exclusively at tight junctions. Claudins are expressed in an organ-specific manner and regulate tissue-specific physiologic properties of tight junctions. By expression in a human intestinal epithelial cell line, it was determined that the intestine-specific homeodomain proteins CDX1 and CDX2 activated a reporter plasmid driven by the CLDN2 promoter.

#### **Application Notes**

Optimal dilution of the Claudin 2 antibody should be determined by the researcher.

#### **Immunogen**

An E. coli-derived human protein (amino acids Q63-V230) was used as the immunogen for the Claudin 2 antibody.

#### **Storage**

After reconstitution, the Claudin 2 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.