
Claudin 2 Antibody / CLDN2 (RQ6463)


Catalog No.	Formulation	Size
RQ6463	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P57739
Applications	Western blot: 1-2ug/ml Immunofluorescence: 5ug/ml Flow cytometry: 1-3ug/million cells Direct ELISA: 0.1-0.5ug/ml
Limitations	This Claudin 2 antibody is available for research use only.



Immunofluorescent staining of FFPE human Caco-2 cells with Claudin 2 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Flow cytometry testing of human A549 cells with Claudin 2 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Claudin 2 antibody.



Western blot testing of human Caco-2 cell lysate with Claudin 2 antibody. Predicted molecular weight ~25 kDa.

Description

Claudin-2 is a protein that in humans is encoded by the CLDN2 gene. By genomic sequence analysis, this gene is mapped to chromosome Xq22.3-q23. This gene product belongs to the claudin protein family whose members have been identified as major integral membrane proteins localized exclusively at tight junctions. Claudins are expressed in an organ-specific manner and regulate tissue-specific physiologic properties of tight junctions. By expression in a human intestinal epithelial cell line, it was determined that the intestine-specific homeodomain proteins CDX1 and CDX2 activated a reporter plasmid driven by the CLDN2 promoter.

Application Notes

Optimal dilution of the Claudin 2 antibody should be determined by the researcher.

Immunogen

An E. coli-derived human protein (amino acids Q63-V230) was used as the immunogen for the Claudin 2 antibody.

Storage

After reconstitution, the Claudin 2 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

Ordering: Phone: 858.663.9055 | Fax: 1.267.821.0800 | Email: info@nsjbio.com
Copyright © NSJ Bioreagents. All rights reserved.