



## Clusterin Antibody / CLU / Apolipoprotein J / APO-J [clone CLU/4730] (V9309)

Catalog No.	Formulation	Size
V9309-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9309-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9309SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


### Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	CLU/4730
Purity	Protein A/G affinity
UniProt	P10909
Localization	Cytoplasm and Nucleus
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This Clusterin antibody is available for research use only.





IHC staining of FFPE human adrenal gland tissue with Clusterin antibody (clone CLU/4730). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Clusterin antibody (clone CLU/4730) as confirmation of integrity and purity.

## Description

Clusterin, also designated complement lysis inhibitor (CLI), apolipoprotein J (APOJ), sulfated glycoprotein 2 (SGP2), SP40 and testosterone-repressed prostate message 2 (TRPM2), is a secretory, heterodimeric glycoprotein that influences immune regulation, cell adhesion, transformation, lipid transportation, tissue remodeling, membrane recycling and cell-cell interactions. Clusterin is synthesized as a 449 amino acid polypeptide that is post-translationally cleaved at an internal bond between Arg 227 and Ser 228. The beta subunit corresponds to residues 23-227. The alpha subunit corresponds to residues 228-449. Overexpression of Clusterin appears to be more common in late stages of mammary tumor progression. Clusterin markedly influences Amyloid structure and neuritic toxicity in vivo and may influence Alzheimer's pathogenesis.

## Application Notes

Optimal dilution of the Clusterin antibody should be determined by the researcher.

## Immunogen

A portion of amino acids 185-287 was used as the immunogen for the Clusterin antibody.

## Storage

Aliquot the Clusterin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.