


Cofilin 2 Antibody / CFL2 [clone CFL2/9963] (V5657)


Catalog No.	Formulation	Size
V5657-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5657-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5657SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	CFL2/9963
Purity	Protein G affinity
UniProt	Q9Y281
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This Cofilin 2 antibody is available for research use only.



IHC staining of FFPE human skeletal muscle tissue with Cofilin 2 antibody (clone CFL2/9963). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Description

Cofilin is ubiquitously expressed in eukaryotic cells where it binds to Actin, thereby regulating the rapid cycling of Actin assembly and disassembly essential for cellular viability. Cofilin is a low molecular weight protein that binds to filamentous (F) Actin by bridging two longitudinally associated Actin subunits changing the F-Actin filament twist. This process is allowed by the dephosphorylation of Cofilin Ser-3 by factors such as opsonized zymosan. Lim kinase 1, a serine kinase, phosphorylates Cofilin and renders it unable to bind and depolymerise F-Actin.

Application Notes

Optimal dilution of the Cofilin 2 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 1-166 from human CFL2 protein was used as the immunogen for the Cofilin 2 antibody.

Storage

Aliquot the Cofilin 2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.