


COPS5 Antibody / JAB1 [clone 4G9] (RQ6537)


Catalog No.	Formulation	Size
RQ6537	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b
Clone Name	4G9
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q92905
Localization	Nuclear and cytoplasmic
Applications	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Immunofluorescence : 5ug/ml Flow Cytometry : 1-3ug/million cells
Limitations	This COPS5 antibody is available for research use only.




IHC staining of FFPE human liver cancer tissue with COPS5 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human placental tissue with COPS5 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human esophageal squamous carcinoma tissue with COPS5 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human lung cancer tissue with COPS5 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE mouse brain tissue with COPS5 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE rat brain tissue with COPS5 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Immunofluorescent staining of FFPE human T-47D cells with COPS5 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of 1) human Raji, 2) human K562, 3) human PC-3, 4) human HeLa, 5) rat brain, 6) rat RH35, 7) mouse brain and 8) mouse SP2/0 cell lysate with COPS5 antibody. Predicted molecular weight ~38 kDa.



Flow cytometry testing of human A549 cells with COPS5 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= COPS5 antibody.

Description

COP9 constitutive photomorphogenic homolog subunit 5 (Arabidopsis), also known as COPS5 or JAB1, is a gene conserved from humans to *Saccharomyces cerevisiae*. It is a member of the MOV34 family. COPS5 is mapped to 8q13.1. The protein encoded by this gene is one of the eight subunits of COP9 signalosome, a highly conserved protein complex that functions as an important regulator in multiple signaling pathways. COPS5 can interact with the cytoplasmic domain of the beta-2 subunit of the alpha-L/beta-2 integrin LFA1, and it is the only protein demonstrated to interact with MIF. COPS5, VHL, and TRC8 proteins appear to be linked both physically and functionally, and all 3 may participate in the development of kidney cancer. In addition to that, COPS5 is an essential cofactor for the apoptotic function of E2F1.

Application Notes

Optimal dilution of the COPS5 antibody should be determined by the researcher.

Immunogen

An E. coli-derived human protein (amino acids M8-S334) was used as the immunogen for the COPS5 antibody.

Storage

After reconstitution, the COPS5 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.