


Creatine kinase B Antibody / CKB [clone 2ba6] (V3497)

Catalog No.	Formulation	Size
V3497-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3497-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3497SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	2ba6
Purity	Protein G affinity chromatography
UniProt	P12277
Localization	Cytoplasmic
Applications	ELISA (order BSA/sodium Azide-free Format For Coating) : Flow Cytometry : 0.5-1ug/10^6 cells Immunofluorescence : 0.5-1ug/ml
Limitations	This Creatine kinase B antibody is available for research use only.



Protein array validation of the Creatine kinase B antibody: Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using Creatine kinase B type antibody (clone 2ba6). These results demonstrate the foremost specificity of the 2ba6 mAb.

Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE Analysis of Purified, BSA-Free Creatine kinase B Antibody (clone 2ba6). Confirmation of Integrity and Purity of the Antibody.

Description

Creatine kinases (CK) are a large family of isoenzymes that regulate levels of ATP in subcellular compartments, where they provide ATP at sites of fluctuating energy demand by the transfer of phosphates between creatine and adenine nucleotides. CKs provide the energy of phosphate hydrolysis necessary to drive the normal function of many cellular systems. In cells, the cytosolic CK enzymes consist of two subunits, which can be either B (brain type) or M (muscle type). There are three different isoenzymes: CKMM, CKBB and CKMB. This mAb recognizes the CKBB isoenzyme and does not react with the B subunit in CKMB. It shows minimal reactivity with other human serum proteins

Application Notes

Optimal dilution of the Creatine kinase B antibody should be determined by the researcher.

Immunogen

Human CKB protein was used as the immunogen for the Creatine kinase B antibody.

Storage

Store the Creatine kinase B antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).