


CSTA Antibody / Cystatin A [clone rCSTA/8995] (V4984)

Catalog No.	Formulation	Size
V4984-100UG	0.2~mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4984-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4984SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Recombinant MOUSE MONOCLONAL

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG2b, kappa
Clone Name	rCSTA/8995
Purity	Protein A/G affinity
UniProt	P01040
Localization	Cell Surface, Cytoplasm, Nucleus
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Limitations	This CSTA antibody is available for research use only.



SDS-PAGE analysis of purified, BSA-free CSTA antibody (clone rCSTA/8995) as confirmation of integrity and purity.

Description

The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins, and kininogens. This gene encodes a stefin that functions as a cysteine protease inhibitor, forming tight complexes with papain and the

cathepsins B, H, and L. The protein is one of the precursor proteins of cornified cell envelope in keratinocytes and plays a role in epidermal development and maintenance. Stefins have been proposed as prognostic and diagnostic tools for cancer.

Application Notes

Optimal dilution of the CSTA antibody should be determined by the researcher.

Immunogen

A recombinant fragment of human protein (around aa 1-98) was used as the immunogen for the CSTA antibody.

Storage

Aliquot the CSTA antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.