


## Cubilin Antibody / CUBN [clone EF-3] (RQ4756)

Catalog No.	Formulation	Size
RQ4756	Antibody in PBS with 0.02% sodium azide, 50% glycerol and 0.4-0.5mg/ml BSA	100 ul

[Bulk quote request](#)

<b>Availability</b>	1-2 weeks
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Rabbit Monoclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Name</b>	EF-3
<b>Purity</b>	Affinity purified
<b>UniProt</b>	O60494
<b>Applications</b>	Western Blot : 1:500-1:2000
<b>Limitations</b>	This Cubilin antibody is available for research use only.



Western blot testing of human fetal kidney with Cubilin antibody. Expected molecular weight: 399-460 kDa.

### Description

Cotransporter which plays a role in lipoprotein, vitamin and iron metabolism, by facilitating their uptake. Binds to ALB, MB, Kappa and lambda-light chains, TF, hemoglobin, GC, SCGB1A1, APOA1, high density lipoprotein, and the CBLIF-cobalamin complex. The binding of all ligands requires calcium. Serves as important transporter in several absorptive epithelia, including intestine, renal proximal tubules and embryonic yolk sac. Interaction with LRP2 mediates its trafficking throughout vesicles and facilitates the uptake of specific ligands like GC, hemoglobin, ALB, TF and SCGB1A1. Interaction with AMN controls its trafficking to the plasma membrane and facilitates endocytosis of ligands. May play an important

role in the development of the peri-implantation embryo through internalization of APOA1 and cholesterol. Binds to LGALS3 at the maternal-fetal interface. [UniProt]

## **Application Notes**

Optimal dilution of the Cubilin antibody should be determined by the researcher.

## **Immunogen**

A synthetic peptide specific to human Cubilin / CUBN was used as the immunogen for the Cubilin antibody.

## **Storage**

Store the Cubilin antibody at -20oC.