CXCL2 Antibody (R31997)

Catalog No.	Formulation	Size
R31997	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	P19875
Applications	Western blot: 0.1-0.5ug/ml ELISA: 0.1-0.5ug/ml (human protein tested); request BSA-free format for coating
Limitations	This CXCL2 antibody is available for research use only.



Western blot testing of human recombinant protein with CXCL2 antibody.

Description

Chemokine (C-X-C motif) ligand 2 is a small cytokine belonging to the CXC chemokine family that is also called macrophage inflammatory protein 2-alpha (MIP2-alpha), Growth-regulated protein beta (Gro-beta) and Gro oncogene-2(Gro-2). CXCL2 is 90% identical in amino acid sequence as a related chemokine, CXCL1. This chemokine is secreted by monocytes and macrophages and is chemotactic for polymorphonuclear leukocytes and hematopoietic stem cells. The gene for CXCL2 is located on human chromosome 4 in a cluster of other CXC chemokines. CXCL2 mobilizes cells by interacting with a cell surface chemokine receptor called CXCR2.

Application Notes

Optimal dilution of the CXCL2 antibody should be determined by the researcher.

Immunogen

Amino acids 39-107 of human CXCL2 were used as the immunogen for the CXCL2 antibody.

Storage

After reconstitution, the CXCL2 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

Copyright © NSJ Bioreagents. All rights reserved.